

Available online at www.sciencedirect.com

ScienceDirect

Procedia - Social and Behavioral Sciences 126 (2014) 168 – 169

Procedia
Social and Behavioral Sciences

ICTTP 2014

Multimodal integration of interval duration: Perceptual or decisional processes?

Maria Dolores de la Rosa*, Karin M. Bausenhardt

Fachbereich Psychologie, University of Tübingen, Schleierstrasse 4, D-72076 Tübingen, Germany

Abstract

Recently, we have shown that perceived duration of visual intervals can be biased by the duration of concurrently presented auditory intervals with conflicting duration (De la Rosa & Bausenhardt, 2013). We suggested that this multimodal biasing effect might be due to temporal ventriloquism (that is, the onset/offset of the visual interval gets ‘pulled’ towards the onset/offset of the auditory interval), which affects the switch component of a pacemaker-accumulator model (Gibbon, 1977). On the other hand, for filled audiovisual intervals with congruent duration, it has been suggested that multimodal integration leads to an increased pacemaker rate of an internal clock (Chen & Yeh, 2009; Walker & Scott, 1981). Importantly, both suggested mechanisms, temporal ventriloquism and pacemaker rate effect, are based on early, perceptual processes. Alternatively, however, it is possible that participants deliberately judge the visual interval duration according to the irrelevant auditory information, and, thus, the observed multimodal integration effects would emerge as the result of a decision process on a relatively late stage of processing. Therefore, in order to distinguish these two mechanisms, we propose an event-related brain potential (ERP) study with the aim to measure the mismatch negativity component (MMN; Näätänen & Alho, 1995). This component mirrors the detection of deviance at an early stage of processing and, thus, indicates underlying pre-attentive and automatic mechanisms. Interestingly, previous studies have already investigated the multimodal integration underlying the spatial ventriloquism effect by assessing the MMN (Stekelenburg, Vroomen, & de Gelder, 2004). It has been found that *illusory* auditory location shifts evoked by real visual location shifts produced an auditory MMN in the auditory cortex. Therefore, it was concluded that the multimodal integration underlying spatial ventriloquism takes place at a pre-attentive level of processing. Similarly, we will investigate the MMN response for conflicting audiovisual interval durations. To this end, our study consists of repeatedly presenting simultaneous audiovisual intervals with a *standard* duration, among which rare stimuli of *deviant* duration are randomly interspersed. Deviants can then occur in: (a) only the auditory modality, (b) only the visual modality, or (c) in both modalities. Additionally, unimodal visual and auditory standards and deviants will be presented as control conditions. We expect that the unimodal auditory and visual deviants evoke an auditory MMN in the auditory cortex and a visual MMN in occipital areas, respectively. Similarly, audiovisual deviants should simultaneously elicit a visual and an auditory MMN in their corresponding areas, probably with an increased magnitude compared to the unimodal conditions. Most interesting are the predictions for conflicting audiovisual intervals: If the multimodal integration of duration is an early perceptual process, auditory-only deviants in bimodal trials should

* Corresponding author. Tel.: +45 (0)7071 29-76423; fax: +49 (0)7071 29-2410.
E-mail address: maria-dolores.de-la-rosa-gamiz@uni-tuebingen.de

bias perceived duration of the visual stimulation, and, thus, a visual MMN in occipital areas should be observed. Because auditory temporal information mostly dominates over visual temporal information, visual-only deviants in bimodal trials should be perceived as similar to the auditory (standard) duration and thus, the visual MMN in occipital areas should even be reduced compared to the unimodal visual MMN. However, if the multimodal integration of perceived duration results from a decision process on a relatively late stage of processing, no MMN for biased perceived duration would be expected, and it should just be evoked by deviant modalities with a real duration shift.

© 2014 The Authors. Published by Elsevier Ltd. Open access under [CC BY-NC-ND license](#).

Selection and peer-review under responsibility of the Organizing Committee of the International Conference on Timing and Time Perception.

Keywords: Time perception; Multisensory integration; Temporal ventriloquism; Mismatch negativity

Acknowledgements

This study was supported by DFG, grant BA 4110/3-1.

References

- Chen, K-M., & Yeh, S-L. (2009). Asymmetric cross-modal effects in time perception. *Acta Psychologica*, 130, 225-234.
- De la Rosa, M. D., & Bausenhardt, K. M. (2013). Multimodal integration of interval duration: Temporal ventriloquism or changes in pacemaker rate? *Timing and Time Perception*, 1, 189-215.
- Gibbon, J. (1977). Scalar expectancy theory and Weber's law in animal timing. *Psychological Review*, 84, 279-325.
- Näätänen, R., & Alho, K. (1995). Mismatch negativity: A unique measure of sensory processing in audition. *International Journal of Neuroscience*, 80, 317- 337.
- Stekelenburg, J. J., Vroomen, J., & de Gelder, B. (2004). Illusory sound shifts induced by the ventriloquist illusion evoke the mismatch negativity. *Neuroscience Letters*, 357, 163-166.
- Walker, J. T., & Scott, K. J. (1981). Auditory-visual conflicts in the perceived duration of lights, tones, and gaps. *Journal of Experimental Psychology Human Perception & Performance*, 7, 1327-1339.